Vector particles tunneling from BTZ black holes

Ge-Rui Chen,* Shiwei Zhou, and Yong-Chang Huang

Institute of Theoretical Physics, Beijing University of Technology, Beijing, 100124, China

Abstract

In this paper we investigate vector particles' Hawking radiation from a BTZ black hole. By applying the WKB approximation and the Hamilton-Jacobi Ansatz to the Proca equation, we obtain the tunneling spectrum of vector particles. The expected Hawking temperature is recovered.

Keywords: Vector particles tunneling, BTZ black hole, Proca equation

arXiv:1409.5941v1 [gr-qc] 21 Sep 2014

*Electronic address: chengerui@emails.bjut.edu.cn

Since Stephen Hawking discovered that black holes radiate black body spectrum[1, 2], there are several different methods to derive Hawking radiation to confirm this astounding discovery. In recent years, semi-classical methods of modeling Hawking radiation as a tunneling process have garnered lots of interest. This semi-classical methods which model Hawking radiation as a tunneling process use WKB approximation to calculate the imaginary part of the action for the classically forbidden trajectory across the horizon. The tunneling probability is given by $\Gamma \propto \exp(-2ImI)$.

The first black hole tunneling method–Null Geodesic Method was proposed by Parikh and Wilczek[3] which followed from the work of Kraus and Wilczek[4]. They found the only part of the action that contributes an imaginary term is $ImI = \int_{r_{in}}^{r_{out}} p_r dr$, where p_r is the momentum of the emitted null s-wave. One can use Hamilton's equation and knowledge of the null geodesics to calculate the imaginary part of the action. Later on, Refs.[5, 6] extended this method to the charged particles. The other method is the Hamilton-Jacobi Ansatz used by Refs.[7, 8], which is an extension of the complex path analysis of Padmanabhan et al[9]. This method applies the WKB approximation to the Klein-Gordon equation, and the lowest order is the Hamilton-Jacobi equation. Then according to the symmetry of the metric, one can pick an appropriate ansatz for the action and put it into the Hamilton-Jacobi equation to solve. In 2008, R. Kerner and R.B. Mann[10, 11] applied the WKB approximation to the Dirac Equation to calculate Dirac particles' Hawking radiation.

Recently, S. I. Kruglov[12] investigated black hole radiation of vector particles in (1+1) dimensions by using the Hamilton-Jacobi method to the Proca equation. Vector particles(e.g. $Z, W^{\pm}$) play important role in Standard Model, so it is interesting to study the Hawking radiation of vector particles. We extend Ref.[12]'s method to investigate vector particles tunneling from a BTZ black hole and obtain the radiation spectrum for vector particles. Our results show that vector particles radiate with the standard Hawking temperature. Like Refs.[10, 11], we assume that the change of black hole angular momentum due to the spin of the emitted particle is negligible. This is a good approximation for the black hole with mass much larger than the Planck mass.

The BTZ back hole is the solution of the standard Einstein-Maxwell equation in $(2+1)$ dimensional spacetime with a negative cosmology constant. For simplicity we will ignore

the coupling to the Maxwell field. The action is

$$I = \frac{1}{2\pi}\int \sqrt{-g}[R + 2l^{-2}]dx^2dt + B, \tag{1}$$

where B is a surface term and the radius l is related to the cosmological constant by $\Lambda = -l^{-2}$. The equations of motion derived from (1) are solved by the black hole field

$$\begin{aligned} ds^2 &= -(-M + \frac{r^2}{l^2} + \frac{J^2}{4r^2})dt^2 + (-M + \frac{r^2}{l^2} + \frac{J^2}{4r^2})^{-1}dr^2 + r^2(d\phi - \frac{J}{2r^2}dt)^2 \\ &= -(-M + \frac{r^2}{l^2})dt^2 + (-M + \frac{r^2}{l^2} + \frac{J^2}{4r^2})^{-1}dr^2 + r^2d\phi^2 - Jdtd\phi \end{aligned} \tag{2}$$

with $-\infty < t < \infty, 0 < r < \infty$ and $0 \leq \phi < 2\pi$. The two constants of integration M and J appearing in (2) are the conserved charges associated with asymptotic invariance under time displacements (mass) and rotational invariance (angular momentum), respectively. The metric (2) has two horizons

$$r_\pm = lM^{\frac{1}{2}}\left[\frac{1}{2}\left(1 \pm \sqrt{1 - (\frac{J}{Ml})^2}\right)\right]^{\frac{1}{2}}, \tag{3}$$

where r_+ is the black hole horizon. In order for the horizon to exist one must have

$$M > 0, |J| \leq Ml. \tag{4}$$

In the extreme case $|J| = Ml$, both roots of g_{11}^{-1} coincide. Let us define the following notations for convenience

$$A(r) = -M + \frac{r^2}{l^2},\ B(r) = -M + \frac{r^2}{l^2} + \frac{J^2}{4r^2},\ C = r^2,\ 2D = -J, \tag{5}$$

so the metric (2) becomes

$$ds^2 = -A(r)dt^2 + \frac{1}{B(r)}dr^2 + Cd\phi^2 + 2Ddtd\phi. \tag{6}$$

The determinant is

$$\sqrt{-g} = \sqrt{\frac{AC + D^2}{B}}, \tag{7}$$

and the inverse metric is

$$g^{\mu\nu} = -\frac{1}{AC + D^2}\begin{pmatrix} C & 0 & -D \\ 0 & -B(AC + D^2) & 0 \\ -D & 0 & -A \end{pmatrix}. \tag{8}$$

The Proca equation for vector particles is[12]

$$D_\mu \psi^{\nu\mu} + \frac{m^2}{\hbar^2}\psi^\nu = 0, \tag{9}$$

$$\psi_{\nu\mu} = D_\nu \psi_\mu - D_\mu \psi_\nu = \partial_\nu \psi_\mu - \partial_\mu \psi_\nu, \tag{10}$$

where D_μ are covariant derivatives, and $\psi_\nu = (\psi_0, \psi_1, \psi_2)$. From the definition, $\psi^{\nu\mu}$ is an anti-symmetrical tensor, so using the equation

$$D_\mu \psi^{\nu\mu} = \frac{1}{\sqrt{-g}}\partial_\mu(\sqrt{-g}\psi^{\nu\mu}), \tag{11}$$

the Proca equation becomes

$$\frac{1}{\sqrt{-g}}\partial_\mu(\sqrt{-g}\psi^{\nu\mu}) + \frac{m^2}{\hbar^2}\psi^\nu = 0. \tag{12}$$

From the metric and the following relationship

$$\begin{aligned}
\psi^0 &= \psi_\mu g^{\mu 0} = \frac{-C}{AC+D^2}\psi_0 + \frac{D}{AC+D^2}\psi_2, \\
\psi^1 &= B\psi_1,\ \psi^2 = \frac{D}{AC+D^2}\psi_0 + \frac{A}{AC+D^2}\psi_2, \\
\psi^{01} &= \psi_{\mu\nu} g^{\mu 0} g^{\nu 1} = -\frac{CB}{AC+D^2}\psi_{01} - \frac{DB}{AC+D^2}\psi_{12}, \\
\psi^{02} &= -\frac{1}{AC+D^2}\psi_{02},\ \psi^{12} = -\frac{BD}{AC+D^2}\psi_{01} + \frac{BA}{AC+D^2}\psi_{12},
\end{aligned} \tag{13}$$

we obtain the Proca equations in the following explicit form

$$\begin{aligned}
&\frac{1}{\sqrt{-g}}\{\partial_r[\sqrt{-g}(\frac{-CB}{AC+D^2}(\partial_t\psi_1 - \partial_r\psi_0) - \frac{DB}{AC+D^2}(\partial_r\psi_2 - \partial_\phi\psi_1))] \\
&+\partial_\phi[\sqrt{-g}(\frac{-1}{AC+D^2}(\partial_t\psi_2 - \partial_\phi\psi_0))]\} + \frac{m^2}{\hbar^2}(\frac{-C}{AC+D^2}\psi_0 + \frac{D}{AC+D^2}\psi_2) = 0, \\
&\frac{1}{\sqrt{-g}}\{\partial_t[\sqrt{-g}(\frac{CB}{AC+D^2}(\partial_t\psi_1 - \partial_r\psi_0) + \frac{DB}{AC+D^2}(\partial_r\psi_2 - \partial_\phi\psi_1))] \\
&+\partial_\phi[\sqrt{-g}(\frac{BD}{AC+D^2}(\partial_r\psi_0 - \partial_t\psi_1) + \frac{AB}{AC+D^2}(\partial_r\psi_2 - \partial_\phi\psi_1))]\} + \frac{m^2}{\hbar^2}B\psi_1 = 0, \\
&\frac{1}{\sqrt{-g}}\{\partial_t[\sqrt{-g}(\frac{1}{AC+D^2}(\partial_t\psi_2 - \partial_\phi\psi_0))] + \partial_r[\sqrt{-g}(\frac{BD}{AC+D^2}(\partial_t\psi_1 - \partial_r\psi_0) \\
&-\frac{AB}{AC+D^2}(\partial_r\psi_2 - \partial_\phi\psi_1))]\} + \frac{m^2}{\hbar^2}(\frac{D}{AC+D^2}\psi_0 + \frac{A}{AC+D^2}\psi_2) = 0.
\end{aligned} \tag{14}$$

According to the WKB approximation, the solutions are in the form

$$\psi_\nu = (c_0, c_1, c_2)\exp(\frac{i}{\hbar}S(t, r, \phi)), \tag{15}$$

where

$$S(t,r,\phi) = S_0(t,r,\phi) + \hbar S_1(t,r,\phi) + \hbar^2 S_2(t,r,\phi) + \cdots \tag{16}$$

Putting Eqs.(15,16) into Eqs.(14), the equations to the leading order in $\hbar$ are

$$\begin{aligned}
&\frac{CB}{AC+D^2}[c_1(\partial_t s_0)(\partial_r s_0) - c_0(\partial_r s_0)^2] + \frac{BD}{AC+D^2}[c_2(\partial_r s_0)^2 - c_1(\partial_\phi s_0)(\partial_r s_0)] \\
&+\frac{1}{AC+D^2}[c_2(\partial_t s_0)(\partial_\phi s_0) - c_0(\partial_\phi s_0)^2] - \frac{C}{AC+D^2}m^2 c_0 + \frac{D}{AC+D^2}m^2 c_2 = 0, \\
&\frac{CB}{AC+D^2}[-c_1(\partial_t s_0)^2 + c_0(\partial_r s_0)(\partial_t s_0)] + \frac{BD}{AC+D^2}[-c_2(\partial_r s_0)(\partial_t s_0) + c_1(\partial_\phi s_0)(\partial_t s_0)] \\
&+\frac{BD}{AC+D^2}[-c_0(\partial_r s_0)(\partial_\phi s_0) + c_1(\partial_t s_0)(\partial_\phi s_0)] \\
&+\frac{AB}{AC+D^2}[-c_2(\partial_r s_0)(\partial_\phi s_0) + c_1(\partial_\phi s_0)^2] + m^2 B c_1 = 0, \\
&\frac{1}{AC+D^2}[-c_2(\partial_t s_0)^2 + c_0(\partial_\phi s_0)(\partial_t s_0)] + \frac{BD}{AC+D^2}[-c_1(\partial_t s_0)(\partial_r s_0) + c_0(\partial_r s_0)^2] \\
&-\frac{AB}{AC+D^2}[-c_2(\partial_r s_0)^2 + c_1(\partial_\phi s_0)(\partial_r s_0)] + m^2\frac{D}{AC+D^2}c_0 + m^2\frac{A}{AC+D^2}c_2 = 0.
\end{aligned} \tag{17}$$

There exists a solution of the form

$$S_0 = -Et + j\phi + W(r) + K, \tag{18}$$

where $E = -\partial_t S_0$ and j are the energy and angular momentum of the emitting particles respectively. K is a complex constant. Inserting Eq.(18) into Eqs.(17) we obtain the matrix equation

$$\Lambda(c_0, c_1, c_2,)^T = 0, \tag{19}$$

where Λ is a 3×3 matrix, and its components are expressed as

$$\begin{aligned}
\Lambda_{11} &= -CB(W')^2 - j^2 - Cm^2,\ \Lambda_{12} = -CBEW' - BDjW', \\
\Lambda_{13} &= BD(W')^2 - Ej + Dm^2,\ \Lambda_{21} = -CBEW' - BDW'j, \\
\Lambda_{22} &= -CBE^2 - 2BDEj + ABj^2 + B(AC+D^2)m^2,\ \Lambda_{23} = BDEW' - ABjW', \\
\Lambda_{31} &= -Ej + BD(W')^2 + m^2 D,\ \Lambda_{32} = BDEW' - BAjW', \\
\Lambda_{33} &= -E^2 + BA(W')^2 + m^2 A,
\end{aligned} \tag{20}$$

where $W' = \partial_r S_0$ and $j = \partial_\phi S_0$.

Homogeneous system of linear equations (19) possesses nontrivial solution if the determinant of the matrix Λ equals to zero, that is, $\det\Lambda = 0$. After the calculation, we have

$$-Bm^2\{-2DEj + Aj^2 + D^2[m^2 + B(W')^2] + C[-E^2 + A(m^2 + B(W')^2)]\}^2 = 0. \quad (21)$$

We obtain immediately

$$\begin{aligned}(W')^2 &= \frac{CE^2 + 2DEj - Aj^2 - m^2(D^2 + CA)}{B(D^2 + CA)} \\ &= \frac{(E - \frac{J}{2r^2}j)^2 - (-M + \frac{r^2}{l^2} + \frac{J^2}{4r^2})(m^2 + \frac{j^2}{r^2})}{(-M + \frac{r^2}{l^2} + \frac{J^2}{4r^2})^2},\end{aligned} \quad (22)$$

and

$$W_\pm = \pm \int \frac{\sqrt{(E - \frac{J}{2r^2}j)^2 - (-M + \frac{r^2}{l^2} + \frac{J^2}{4r^2})(m^2 + \frac{j^2}{r^2})}}{-M + \frac{r^2}{l^2} + \frac{J^2}{4r^2}} dr. \quad (23)$$

One solution W_+ corresponds to vector particles moving away from the black hole, and the other W_- corresponds to particles moving toward the black hole. Based on the discussion of Ref.[10], the probability of tunneling particles is

$$\Gamma \propto \exp(-\frac{4}{\hbar} Im W_+). \quad (24)$$

Integrating around the pole at the horizon r_+, we obtain

$$Im W_+ = \frac{\pi}{2\kappa}(E - \Omega_+ j), \quad (25)$$

where $\kappa = \frac{r_+^2 - r_-^2}{l^2 r_+}$ is the surface gravity of outer event horizon and $\Omega_+ \equiv -\frac{g_{02}}{g_{22}}|_{r_+} = \frac{J}{2r_+^2}$ is the angular velocity of the outer event horizon. So the tunneling probability is

$$\Gamma \propto \exp[-\frac{2\pi}{\kappa}(E - \Omega_+ j)], \quad (26)$$

and the Hawking temperature is

$$T_H = \frac{\kappa}{2\pi} = \frac{r_+^2 - r_-^2}{2\pi l^2 r_+}. \quad (27)$$

We get the Hawking temperature of vector particles from BTZ black holes. The result is the same as Dirac particles tunneling from BTZ black holes[13].

In summary, we investigate vector particles' Hawking radiation from a BTZ black hole. By applying the WKB approximation and the Hamilton-Jacobi Ansatz to the Proca equation, we obtain the tunneling spectrum of vector particles and the expected Hawking temperature, which provides further evidence for the universality of back radiation. Compared with other methods to calculate Hawking radiation, Hamilton-Jacobi method has a clear physical picture, is straightforward to compute, and easily generalize to different kinds of particles.

Acknowledgments

This work is supported by National Natural Science Foundation of China (No.11275017 and No.11173028).

---

[1] S. Hawking, Nature **248**(1974)30.

[2] S. Hawking, Commun. Math. Phys. **43**(1975)199.

[3] M. K. Parikh, F.Wilczek, Phys. Rev. Lett. **85**(2000)5042; M. K. Parikh, Int. J. Mod. Phys. D **13**(2004)2351.

[4] P. Kraus and F. Wilczek, Nucl. Phys. B **433**(1995)403; P. Kraus and F. Wilczek, Nucl. Phys. B **437**(1995)231-242; P. Kraus and E. Keski-Vakkuri, Nucl.Phys. B **491**(1997)249-262.

[5] Qing-Quan Jiang, Shuang-Qing Wu, Xu Cai, Phys.Rev. D **73**(2006)064003.

[6] J. Zhang and Z. Zhao, Phys. Lett. B **638**(2006)110.

[7] M. Angheben, M. Nadalini, L. Vanzo, and S. Zerbini, J. High Energy Phys. **05**(2005)014.

[8] R. Kerner and R. B. Mann, Phys. Rev. D **73**(2006)104010.

[9] K. Srinivasan and T. Padmanabhan, Phys. Rev. D **60**(1999)24007; S. Shankaranarayanan, K. S rinivasan, and T. Padmanabhan, Mod.Phys.Lett. A **16**(2001)571-578; S. Shankaranarayanan, T. Padmanabhan, and K. Srinivasan, Class.Quant.Grav. **19**(2002)2671-2688; S. Shankaranarayanan, Phys.Rev. D **67**(2003)084026.

[10] R. Kerner and R. B. Mann, Class. Quant. Grav. **25**(2008)095014.

[11] R. Kerner and R. B. Mann, Phys. Lett. B **665**(2008)277-283.

[12] S. I. Kruglov, Int. J. Mod. Phys. A **29**(2014)1450118.

[13] Ran Li, Ji-Rong Ren, Phys.Lett.B **661**(2008)370-372.